

ICM7068 - The Physiology of Shock, Shock Syndromes and Tools of Resuscitation

View Online



This module is designed to teach the student how to identify which patients require resuscitation and the tools by which this is achieved.

1

Guyton AC, Hall JE. Guyton and Hall textbook of medical physiology. Thirteenth edition. Philadelphia, PA: : Elsevier 2016.
<https://www-vlebooks-com.ezproxy.library.qmul.ac.uk/Vleweb/Product/Index/748473?page=0>

2

Tuma M, Canestrini S, Alwahab Z, et al. Trauma and Endothelial Glycocalyx. SHOCK 2016; **46**:352-7. doi:10.1097/SHK.0000000000000635

3

Piehl MD, Manning JE, McCurdy SL, et al. Pulse contour cardiac output analysis in a piglet model of severe hemorrhagic shock*. Critical Care Medicine 2008;**36**:1189-95.
doi:10.1097/CCM.0B013E31816592A3

4

American Journal of Physiology-Regulatory, Integrative and Comparative Physiology.
<https://www.physiology.org/doi/full/10.1152/ajpregu.00304.2015>

5

Lord JM, Midwinter MJ, Chen Y-F, et al. The systemic immune response to trauma: an overview of pathophysiology and treatment. The Lancet 2014;**384**:1455-65.
doi:10.1016/S0140-6736(14)60687-5

6

Brohi K, Cohen MJ, Davenport RA. Acute coagulopathy of trauma: mechanism, identification and effect. *Current Opinion in Critical Care* 2007;**13**:680-5.
doi:10.1097/MCC.0b013e3282f1e78f

7

Pranskunas A, Koopmans M, Koetsier PM, et al. Microcirculatory blood flow as a tool to select ICU patients eligible for fluid therapy. *Intensive Care Medicine* 2013;**39**:612-9.
doi:10.1007/s00134-012-2793-8

8

Trzeciak S, Dellinger RP, Parrillo JE, et al. Early microcirculatory perfusion derangements in patients with severe sepsis and septic shock: Relationship to hemodynamics, oxygen transport, and survival. *Annals of Emergency Medicine* 2007;**49**:88-98.e2.
doi:10.1016/j.annemergmed.2006.08.021

9

Jhanji S, Lee C, Watson D, et al. Microvascular flow and tissue oxygenation after major abdominal surgery: association with post-operative complications. *Intensive Care Medicine* 2009;**35**:671-7. doi:10.1007/s00134-008-1325-z

10

Tachon G, Harrois A, Tanaka S, et al. Microcirculatory Alterations in Traumatic Hemorrhagic Shock*. *Critical Care Medicine* 2014;**42**:1433-41.
doi:10.1097/CCM.0000000000000223

11

Hutchings SD, Naumann DN, Watts S, et al. Microcirculatory perfusion shows wide inter-individual variation and is important in determining shock reversal during resuscitation in a porcine experimental model of complex traumatic hemorrhagic shock. *Intensive Care Medicine Experimental* 2016;**4**. doi:10.1186/s40635-016-0088-z

12

Jhanji S, Stirling S, Patel N, et al. The effect of increasing doses of norepinephrine on tissue oxygenation and microvascular flow in patients with septic shock*. *Critical Care Medicine* 2009;**37**:1961–6. doi:10.1097/CCM.0b013e3181a00a1c

13

Morelli A, Donati A, Ertmer C, et al. Effects of vasopressinergic receptor agonists on sublingual microcirculation in norepinephrine-dependent septic shock. *Critical Care* 2011;**15**. doi:10.1186/cc10453

14

Morelli A, Donati A, Ertmer C, et al. Levosimendan for resuscitating the microcirculation in patients with septic shock: a randomized controlled study. *Critical Care* 2010;**14**. doi:10.1186/cc9387

15

Jhanji S, Vivian-Smith A, Lucena-Amaro S, et al. Haemodynamic optimisation improves tissue microvascular flow and oxygenation after major surgery: a randomised controlled trial. *Critical Care* 2010;**14**. doi:10.1186/cc9220

16

Xu J, Ma L, Sun S, et al. Fluid Resuscitation Guided by Sublingual Partial Pressure of Carbon Dioxide During Hemorrhagic Shock in a Porcine Model. *Shock* 2013;**39**:361–5. doi:10.1097/SHK.0b013e31828936aa

17

Shoemaker WC, Appel PL, Kram HB, et al. Prospective Trial of Supranormal Values of Survivors as Therapeutic Goals in High-Risk Surgical Patients. *Chest* 1988;**94**:1176–86. doi:10.1378/chest.94.6.1176

18

Starodub R, Abella BS, Grossestreuer AV, et al. Association of serum lactate and survival

outcomes in patients undergoing therapeutic hypothermia after cardiac arrest. Resuscitation 2013;**84**:1078–82. doi:10.1016/j.resuscitation.2013.02.001

19

Marik PE, Baram M, Vahid B. Does Central Venous Pressure Predict Fluid Responsiveness?*: A Systematic Review of the Literature and the Tale of Seven Mares. Chest 2008;**134**:172–8. doi:10.1378/chest.07-2331

20

Pierrakos C, Velissaris D, Scolletta S, et al. Can changes in arterial pressure be used to detect changes in cardiac index during fluid challenge in patients with septic shock? Intensive Care Medicine 2012;**38**:422–8. doi:10.1007/s00134-011-2457-0

21

Alhashemi JA, Cecconi M, Hofer CK. Cardiac output monitoring: an integrative perspective. Critical Care 2011;**15**. doi:10.1186/cc9996

22

Cecconi M, Hofer C, Teboul J-L, et al. Fluid challenges in intensive care: the FENICE study. Intensive Care Medicine 2015;**41**:1529–37. doi:10.1007/s00134-015-3850-x

23

Pierrakos C, Velissaris D, Scolletta S, et al. Can changes in arterial pressure be used to detect changes in cardiac index during fluid challenge in patients with septic shock? Intensive Care Medicine 2012;**38**:422–8. doi:10.1007/s00134-011-2457-0

24

Monge García MI, Guijo González P, Gracia Romero M, et al. Effects of fluid administration on arterial load in septic shock patients. Intensive Care Medicine 2015;**41**:1247–55. doi:10.1007/s00134-015-3898-7

25

Aya HD, Ster IC, Fletcher N, et al. Pharmacodynamic Analysis of a Fluid Challenge. *Critical Care Medicine* 2016;**44**:880–91. doi:10.1097/CCM.0000000000001517

26

Aya HD, Rhodes A, Chis Ster I, et al. Hemodynamic Effect of Different Doses of Fluids for a Fluid Challenge. *Critical Care Medicine* 2017;**45**:e161–8. doi:10.1097/CCM.0000000000002067

27

Nunes TSO, Ladeira RT, Bafi AT, et al. Duration of hemodynamic effects of crystalloids in patients with circulatory shock after initial resuscitation. *Annals of Intensive Care* 2014;**4**. doi:10.1186/s13613-014-0025-9

28

Monnet X, Rienzo M, Osman D, et al. Passive leg raising predicts fluid responsiveness in the critically ill*. *Critical Care Medicine* 2006;**34**:1402–7. doi:10.1097/01.CCM.0000215453.11735.06

29

Clarke DL, Chipps JA, Sartorius B, et al. Mortality rates increase dramatically below a systolic blood pressure of 105-mm Hg in septic surgical patients. *The American Journal of Surgery* 2016;**212**:941–5. doi:10.1016/j.amjsurg.2016.01.042

30

Lee YK, Hwang SY, Shin TG, et al. Prognostic Value of Lactate and Central Venous Oxygen Saturation after Early Resuscitation in Sepsis Patients. *PLOS ONE* 2016;**11**. doi:10.1371/journal.pone.0153305

31

Angus DC, van der Poll T. Severe Sepsis and Septic Shock. *New England Journal of Medicine* 2013;**369**:840–51. doi:10.1056/NEJMra1208623

32

A Randomized Trial of Protocol-Based Care for Early Septic Shock. *The New England Journal of Medicine*; **370**:1683–93. <https://search.proquest.com/docview/1520423050?pq-origsite=summon>

33

Mellhammar L, Wullt S, Lindberg Å, et al. Sepsis Incidence: A Population-Based Study. *Open Forum Infectious Diseases* 2016;**3**. doi:10.1093/ofid/ofw207

34

Assessment of Global Incidence and Mortality of Hospital-treated Sepsis. Current Estimates and Limitations | *American Journal of Respiratory and Critical Care Medicine*. <https://www.atsjournals.org/doi/10.1164/rccm.201504-0781OC>

35

Rhodes A, Evans LE, Alhazzani W, et al. Surviving Sepsis Campaign. *Critical Care Medicine* 2017;**45**:486–552. doi:10.1097/CCM.0000000000002255

36

Reinhart K, Daniels R, Kisson N, et al. Recognizing Sepsis as a Global Health Priority — A WHO Resolution. *New England Journal of Medicine* 2017;**377**:414–7. doi:10.1056/NEJMp1707170

37

Tackling drug-resistant infections globally. https://amr-review.org/sites/default/files/160518_Final%20paper_with%20cover.pdf

38

A European One Health Action Plan against Antimicrobial Resistance. https://ec.europa.eu/health/sites/health/files/antimicrobial_resistance/docs/amr_2017_actio

n-plan.pdf

39

De Backer D, Cecconi M, Lipman J, et al. Challenges in the management of septic shock: a narrative review. *Intensive Care Medicine* Published Online First: 11 February 2019. doi:10.1007/s00134-019-05544-x

40

Reynolds HR, Hochman JS. Cardiogenic Shock: Current Concepts and Improving Outcomes. *Circulation* 2008;**117**:686–97. doi:10.1161/CIRCULATIONAHA.106.613596

41

Haemodynamic Monitoring: ESICM EDIC PACT Study Tool.
<https://www.esicm.org/education/>

42

Elliott P. Rational use of inotropes. *Anaesthesia & Intensive Care Medicine* 2006;**7**:326–30. doi:10.1053/j.mpaic.2006.06.007

43

Napp LC, Kühn C, Bauersachs J. ECMO in cardiac arrest and cardiogenic shock. *Herz* 2017;**42**:27–44. doi:10.1007/s00059-016-4523-4

44

Mebazaa A, Nieminen MS, Packer M, et al. Levosimendan vs Dobutamine for Patients With Acute Decompensated Heart Failure. *JAMA* 2007;**297**. doi:10.1001/jama.297.17.1883

45

Gray A, Goodacre S, Newby DE, et al. Noninvasive Ventilation in Acute Cardiogenic Pulmonary Edema. *New England Journal of Medicine* 2008;**359**:142–51.

doi:10.1056/NEJMoa0707992

46

Masip J, Roque M, Sánchez B, et al. Noninvasive Ventilation in Acute Cardiogenic Pulmonary Edema. *JAMA* 2005;**294**. doi:10.1001/jama.294.24.3124

47

Sackner-Bernstein JD, Kowalski M, Fox M, et al. Short-term Risk of Death After Treatment With Nesiritide for Decompensated Heart Failure. *JAMA* 2005;**293**. doi:10.1001/jama.293.15.1900

48

Thiele H, Zeymer U, Neumann F-J, et al. Intraaortic Balloon Support for Myocardial Infarction with Cardiogenic Shock. *New England Journal of Medicine* 2012;**367**:1287–96. doi:10.1056/NEJMoa1208410

49

Thiele H, Jobs A, Ouweneel DM, et al. Percutaneous short-term active mechanical support devices in cardiogenic shock: a systematic review and collaborative meta-analysis of randomized trials. *European Heart Journal* 2017;**38**:3523–31. doi:10.1093/eurheartj/ehx363

50

Circulation. <https://www.ahajournals.org/doi/10.1161/CIR.0000000000000509>

51

Booth RA, Hill SA, Don-Wauchope A, et al. Performance of BNP and NT-proBNP for diagnosis of heart failure in primary care patients: a systematic review. *Heart Failure Reviews* 2014;**19**:439–51. doi:10.1007/s10741-014-9445-8

52

Napp LC, Kühn C, Bauersachs J. ECMO in cardiac arrest and cardiogenic shock. *Herz* 2017;
42:27-44. doi:10.1007/s00059-016-4523-4

53

Major trauma: assessment and initial management | Guidance and guidelines | NICE.
<https://www.nice.org.uk/guidance/ng39>

54

Acute upper gastrointestinal bleeding in over 16s: management | Guidance and guidelines
| NICE. <https://www.nice.org.uk/guidance/cg141/chapter/1-guidance>

55

The European Society of GI Endoscopy (ESGE) Guideline on the diagnosis and
management of nonvariceal UGI haemorrhage.
https://www.esge.com/assets/downloads/pdfs/guidelines/2015_s_0034_1393172.pdf

56

Solomon, Caren GLaine, Loren. Upper Gastrointestinal Bleeding Due to a Peptic Ulcer. *The
New England Journal of Medicine*; **374**
:2367-76. <https://search.proquest.com/docview/1798243079?pq-origsite=summon>

57

Laine, Loren, MD. Blood Transfusion for Gastrointestinal Bleeding. *The New England Journal
of Medicine*; **368**
:75-6. <https://search.proquest.com/docview/1266235512?pq-origsite=summon>

58

Oyeniya BT, Fox EE, Scerbo M, et al. Trends in 1029 trauma deaths at a level 1 trauma
center: Impact of a bleeding control bundle of care. *Injury* 2017; **48**:5-12.
doi:10.1016/j.injury.2016.10.037

59

Holcomb JB, Tilley BC, Baraniuk S, et al. Transfusion of Plasma, Platelets, and Red Blood Cells in a 1:1:1 vs a 1:1:2 Ratio and Mortality in Patients With Severe Trauma. *JAMA* 2015; **313**. doi:10.1001/jama.2015.12

60

Cannon JW, Khan MA, Raja AS, et al. Damage control resuscitation in patients with severe traumatic hemorrhage. *Journal of Trauma and Acute Care Surgery* 2017;**82**:605–17. doi:10.1097/TA.0000000000001333

61

Shackelford SA, del Junco DJ, Powell-Dunford N, et al. Association of Prehospital Blood Product Transfusion During Medical Evacuation of Combat Casualties in Afghanistan With Acute and 30-Day Survival. *JAMA* 2017;**318**. doi:10.1001/jama.2017.15097

62

Spinella PC, Pidcoke HF, Strandenes G, et al. Whole blood for hemostatic resuscitation of major bleeding. *Transfusion* 2016;**56**:S190–202. doi:10.1111/trf.13491

63

Holcomb JB. Reconstitution: Reverse Engineering. *The Journal of Trauma: Injury, Infection, and Critical Care* 2011;**70**:S65–7. doi:10.1097/TA.0b013e31821a609c

64

Glassberg E, Nadler R, Gendler S, et al. Freeze-Dried Plasma at the Point of Injury. *Shock* 2013;**40**:444–50. doi:10.1097/SHK.0000000000000047

65

An Ethical Framework for Controlled Donation after Circulatory Death: Executive Summary - Academy of Medical Royal Colleges.
<http://www.aomrc.org.uk/publications/reports-guidance/ethical-framework-controlled-donation-circulatory-death-executive-summary/>

66

A code of practice for the diagnosis and confirmation of death - Academy of Medical Royal Colleges.

<http://www.aomrc.org.uk/publications/reports-guidance/code-practice-diagnosis-confirmation-death/>

67

Blackstock MJ, Ray DC. Organ donation after circulatory death. *European Journal of Emergency Medicine* 2014;**21**:324–9. doi:10.1097/MEJ.0000000000000082

68

Sampson, Hugh A Muñoz-Furlong, Anne Campbell, Ronna L Adkinson, N Franklin Bock, S Allan. Second symposium on the definition and management of anaphylaxis: Summary report--Second National Institute of Allergy and Infectious Disease/Food Allergy and Anaphylaxis Network symposium. *Journal of Allergy and Clinical Immunology*; **117**:391–7. <https://search.proquest.com/docview/1504744658?pq-origsite=summon>

69

Jiwaji Z, Brady S, McIntyre LA, et al. Emergency department management of early sepsis: a national survey of emergency medicine and intensive care consultants. *Emergency Medicine Journal* 2014;**31**:1000–5. doi:10.1136/emmermed-2013-202883

70

A Comparison of Albumin and Saline for Fluid Resuscitation in the Intensive Care Unit. *New England Journal of Medicine* 2004;**350**:2247–56. doi:10.1056/NEJMoa040232

71

Maitland K, Kiguli S, Opoka RO, et al. Mortality after Fluid Bolus in African Children with Severe Infection. *New England Journal of Medicine* 2011;**364**:2483–95. doi:10.1056/NEJMoa1101549

72

Myburgh JA, Finfer S, Bellomo R, et al. Hydroxyethyl Starch or Saline for Fluid Resuscitation in Intensive Care. *New England Journal of Medicine* 2012;**367**:1901–11.
doi:10.1056/NEJMoa1209759

73

Antonelli M, Sandroni C. Hydroxyethyl Starch for Intravenous Volume Replacement. *JAMA* 2013;**309**. doi:10.1001/jama.2013.851

74

Young P, Bailey M, Beasley R, et al. Effect of a Buffered Crystalloid Solution vs Saline on Acute Kidney Injury Among Patients in the Intensive Care Unit. *JAMA* 2015;**314**.
doi:10.1001/jama.2015.12334

75

Vincent JL. Anemia and Blood Transfusion in Critically Ill Patients. *JAMA* 2002;**288**.
doi:10.1001/jama.288.12.1499

76

Rodriguez RM, Lum-Lung M, Dixon K, et al. A prospective study on esophageal Doppler hemodynamic assessment in the ED. *The American Journal of Emergency Medicine* 2006;
24:658–63. doi:10.1016/j.ajem.2006.02.006

77

Howell MD, Donnino M, Clardy P, et al. Occult hypoperfusion and mortality in patients with suspected infection. *Intensive Care Medicine* 2007;**33**:1892–9.
doi:10.1007/s00134-007-0680-5

78

Helmerhorst HJF, Roos-Blom M-J, van Westerloo DJ, et al. Association Between Arterial Hyperoxia and Outcome in Subsets of Critical Illness. *Critical Care Medicine* 2015;**43**:1508–19. doi:10.1097/CCM.0000000000000998

79

Kilgannon JH. Association Between Arterial Hyperoxia Following Resuscitation From Cardiac Arrest and In-Hospital Mortality. *JAMA* 2010;**303**. doi:10.1001/jama.2010.707

80

Stub D, Smith K, Bernard S, et al. Air Versus Oxygen in ST-Segment-Elevation Myocardial Infarction. *Circulation* 2015;**131**:2143–50. doi:10.1161/CIRCULATIONAHA.114.014494

81

Rincon F, Kang J, Maltenfort M, et al. Association Between Hyperoxia and Mortality After Stroke. *Critical Care Medicine* 2014;**42**:387–96. doi:10.1097/CCM.0b013e3182a27732

82

Perel P, Roberts I, Ker K. Colloids versus crystalloids for fluid resuscitation in critically ill patients. *Cochrane Database of Systematic Reviews Published Online First*: 28 February 2013. doi:10.1002/14651858.CD000567.pub6

83

Young P, Bailey M, Beasley R, et al. Effect of a Buffered Crystalloid Solution vs Saline on Acute Kidney Injury Among Patients in the Intensive Care Unit. *JAMA* 2015;**314**. doi:10.1001/jama.2015.12334

84

Angus DC, Barnato AE, Bell D, et al. A systematic review and meta-analysis of early goal-directed therapy for septic shock: the ARISE, ProCESS and ProMISe Investigators. *Intensive Care Medicine* 2015;**41**:1549–60. doi:10.1007/s00134-015-3822-1

85

Maitland K, Kiguli S, Opoka RO, et al. Mortality after Fluid Bolus in African Children with Severe Infection. *New England Journal of Medicine* 2011;**364**:2483–95.

doi:10.1056/NEJMoa1101549

86

Maitland K, George EC, Evans JA, et al. Exploring mechanisms of excess mortality with early fluid resuscitation: insights from the FEAST trial. *BMC Medicine* 2013;**11**.
doi:10.1186/1741-7015-11-68

87

Ospina-Tascon G, Neves AP, Occhipinti G, et al. Effects of fluids on microvascular perfusion in patients with severe sepsis. *Intensive Care Medicine* 2010;**36**:949-55.
doi:10.1007/s00134-010-1843-3

88

Glassford N, Eastwood G, Bellomo R. Physiological changes after fluid bolus therapy in sepsis: a systematic review of the contemporary literature. *Critical Care* 2014;**18**.
doi:10.1186/cc14037

89

Guidet B, Ait-Oufella H. Fluid resuscitation should respect the endothelial glycocalyx layer. *Critical Care* 2014;**18**. doi:10.1186/s13054-014-0707-6

90

Chappell D, Bruegger D, Potzel J, et al. Hypervolemia increases release of atrial natriuretic peptide and shedding of the endothelial glycocalyx. *Critical Care* 2014;**18**.
doi:10.1186/s13054-014-0538-5

91

Young PP, Cotton BA, Goodnough LT. Massive Transfusion Protocols for Patients With Substantial Hemorrhage. *Transfusion Medicine Reviews* 2011;**25**:293-303.
doi:10.1016/j.tmr.2011.04.002

92

Hogshire L, Carson JL. Red blood cell transfusion. Current Opinion in Hematology 2013;**20**:546-51. doi:10.1097/MOH.0b013e32836508bd

93

Lucas CE, Ledgerwood AM. FFP:RBC Resuscitation Ratio and Post-Shock Fluid Uptake. JAMA Surgery 2013;**148**:623. doi:10.1001/jamasurg.2013.623

94

Gonzalez EA, Moore FA, Holcomb JB, et al. Fresh Frozen Plasma Should be Given Earlier to Patients Requiring Massive Transfusion. The Journal of Trauma: Injury, Infection, and Critical Care 2007;**62**:112-9. doi:10.1097/01.ta.0000250497.08101.8b

95

Guidelines | British Society for Haematology. <http://www.b-s-h.org.uk/guidelines/>

96

Association of Anaesthetists of Great Britain and Ireland. <http://www.aagbi.org/>

97

Handbook of Transfusion Medicine.
<https://www.transfusionguidelines.org/transfusion-handbook>

98

EMCrit Blog - Emergency Department Critical Care & Resuscitation. <https://emcrit.org/>

99

Nevin DG, Brohi K. Permissive hypotension for active haemorrhage in trauma. Anaesthesia Published Online First: 22 September 2017. doi:10.1111/anae.14034

100

Myles PS, Smith JA, Forbes A, et al. Tranexamic Acid in Patients Undergoing Coronary-Artery Surgery. *New England Journal of Medicine* 2017;**376**:136–48. doi:10.1056/NEJMoa1606424

101

Gayet-Ageron A, Prieto-Merino D, Ker K, et al. Effect of treatment delay on the effectiveness and safety of antifibrinolytics in acute severe haemorrhage: a meta-analysis of individual patient-level data from 40 138 bleeding patients. *The Lancet* 2018;**391**:125–32. doi:10.1016/S0140-6736(17)32455-8

102

World Population Ageing 2015.
http://www.un.org/en/development/desa/population/publications/pdf/ageing/WPA2015_Report.pdf

103

Physiology of Ageing.
<https://www.sciencedirect.com/science/article/abs/pii/S1357303916302298>

104

Sieck GC. Physiology of aging. *Journal of Applied Physiology* 2003;**95**:1333–4. doi:10.1152/jappphysiol.00718.2003

105

Sammy I, Lecky F, Sutton A, et al. Factors affecting mortality in older trauma patients—A systematic review and meta-analysis. *Injury* 2016;**47**:1170–83. doi:10.1016/j.injury.2016.02.027

106

Cook I, Kirkup AL, Langham LJ, et al. End of Life Care and Do Not Resuscitate Orders: How Much Does Age Influence Decision Making? A Systematic Review and Meta-Analysis. *Gerontology and Geriatric Medicine* 2017;**3**. doi:10.1177/2333721417713422

107

O'Grady NP, Alexander M, Burns LA, et al. Guidelines for the prevention of intravascular catheter-related infections. *American Journal of Infection Control* 2011;**39**:S1-34. doi:10.1016/j.ajic.2011.01.003

108

Prytherch DR, Smith GB, Schmidt PE, et al. ViEWS—Towards a national early warning score for detecting adult inpatient deterioration. *Resuscitation* 2010;**81**:932-7. doi:10.1016/j.resuscitation.2010.04.014

109

Burch VC, Tarr G, Morroni C. Modified early warning score predicts the need for hospital admission and inhospital mortality. *Emergency Medicine Journal* 2008;**25**:674-8. doi:10.1136/emj.2007.057661