

ICM7062 - Trauma to the Pelvis and Torso

[View Online](#)

[1]

Adler, E. 2011. Defining the Limits of Resuscitative Emergency Department Thoracotomy: A Contemporary Western Trauma Association Perspective. *The Journal of Emergency Medicine*. 41, 2 (Aug. 2011), 231-232.

DOI:<https://doi.org/10.1016/j.jemermed.2011.06.028>.

[2]

Blunt Hepatic Injury, Selective Nonoperative Management of - Practice Management Guideline:

<https://www.east.org/education-resources/practice-management-guidelines/details/hepatic-injury-blunt-selective-nonoperative-management-of>.

[3]

Burlew, C.C. 2017. Preperitoneal pelvic packing for exsanguinating pelvic fractures. *International Orthopaedics*. 41, 9 (Sep. 2017), 1825-1829.

DOI:<https://doi.org/10.1007/s00264-017-3485-3>.

[4]

Cheatham, M.L. et al. 2007. Results from the International Conference of Experts on Intra-abdominal Hypertension and Abdominal Compartment Syndrome. II. Recommendations. *Intensive Care Medicine*. 33, 6 (May 2007), 951-962.

DOI:<https://doi.org/10.1007/s00134-007-0592-4>.

[5]

Crichton, J.C.I. et al. 2017. The role of splenic angioembolization as an adjunct to

nonoperative management of blunt splenic injuries. *Journal of Trauma and Acute Care Surgery*. 83, 5 (Nov. 2017), 934–943. DOI:<https://doi.org/10.1097/TA.0000000000001649>.

[6]

Demetriades, D. et al. 2002. Pelvic fractures: epidemiology and predictors of associated abdominal injuries and outcomes¹ 1No competing interests declared. *Journal of the American College of Surgeons*. 195, 1 (Jul. 2002), 1–10. DOI:[https://doi.org/10.1016/S1072-7515\(02\)01197-3](https://doi.org/10.1016/S1072-7515(02)01197-3).

[7]

Filiberto, D.M. and Fox, A.D. 2016. Preperitoneal pelvic packing: Technique and outcomes. *International Journal of Surgery*. 33, (Sep. 2016), 222–224. DOI:<https://doi.org/10.1016/j.ijssu.2016.05.072>.

[8]

Gustavo Parreira, J. et al. 2000. The role of associated injuries on outcome of blunt trauma patients sustaining pelvic fractures. *Injury*. 31, 9 (Nov. 2000), 677–682. DOI:[https://doi.org/10.1016/S0020-1383\(00\)00074-7](https://doi.org/10.1016/S0020-1383(00)00074-7).

[9]

Huh, J. et al. 2003. Surgical management of traumatic pulmonary injury. *The American Journal of Surgery*. 186, 6 (Dec. 2003), 620–624. DOI:<https://doi.org/10.1016/j.amjsurg.2003.08.013>.

[10]

Kozar, R.A. 2006. Risk Factors for Hepatic Morbidity Following Nonoperative Management. *Archives of Surgery*. 141, 5 (May 2006). DOI:<https://doi.org/10.1001/archsurg.141.5.451>.

[11]

Kozar, R.A. and McNutt, M.K. 2010. Management of adult blunt hepatic trauma. *Current Opinion in Critical Care*. 16, 6 (Dec. 2010), 596–601.

DOI:<https://doi.org/10.1097/MCC.0b013e32833f5cd5>.

[12]

Mattox, Kenneth L. et al. 2013. Trauma. McGraw-Hill Medical.

[13]

Noser, J. et al. 2018. Mid-term follow-up after surgical treatment of fragility fractures of the pelvis. *Injury*. 49, 11 (Nov. 2018), 2032–2035.
DOI:<https://doi.org/10.1016/j.injury.2018.09.017>.

[14]

Pancreatic Trauma, Diagnosis and Management of - Practice Management Guideline:
<https://www.east.org/education-resources/practice-management-guidelines/archived/pancreatic-trauma-diagnosis-and-management-of>.

[15]

Patterson, L 2002. Practice management guidelines for the evaluation of blunt abdominal trauma: The EAST practice management guidelines work group. *JOURNAL OF TRAUMA-INJURY INFECTION AND CRITICAL CARE*. 53, 3 (Sep. 2002), 602–615.
DOI:<https://doi.org/10.1097/01.TA.0000025413.43206.97>.

[16]

Sathy, A.K. et al. 2009. The Effect of Pelvic Fracture on Mortality After Trauma: An Analysis of 63,000 Trauma Patients. *The Journal of Bone and Joint Surgery-American Volume*. 91, 12 (Dec. 2009), 2803–2810. DOI:<https://doi.org/10.2106/JBJS.H.00598>.

[17]

Shrestha, B. et al. 2015. Damage-control resuscitation increases successful nonoperative management rates and survival after severe blunt liver injury. *Journal of Trauma and Acute Care Surgery*. 78, 2 (Feb. 2015), 336–341.
DOI:<https://doi.org/10.1097/TA.0000000000000514>.

[18]

The EAST approach to guideline development using GRADE methodology:
<http://www.east.org/resources/treatment-guidelines>.

[19]

Wiley, M. et al. 2018. Complications After Pelvic Arteriography in Patients With Pelvic Ring Disruptions. *Journal of the American Academy of Orthopaedic Surgeons*. 26, 21 (Nov. 2018), 765–772. DOI:<https://doi.org/10.5435/JAAOS-D-17-00198>.

[20]

Zarzaaur, B.L. et al. 2017. Natural history of splenic vascular abnormalities after blunt injury. *Journal of Trauma and Acute Care Surgery*. 83, 6 (Dec. 2017), 999–1005. DOI:<https://doi.org/10.1097/TA.0000000000001597>.

[21]

2001. Practice management guidelines for prophylactic antibiotic use in penetrating abdominal trauma: The EAST Practice management guidelines work group. *The American Journal of Emergency Medicine*. 19, 5 (Sep. 2001). DOI:[https://doi.org/10.1016/S0735-6757\(01\)80146-7](https://doi.org/10.1016/S0735-6757(01)80146-7).

[22]

The Bone & Joint Journal.